


EXTENDED COANNIHILATIONS FROM NON UNIVERSAL SFERMION MASSES

S. PROFUMO

*Scuola Internazionale Superiore di Studi Avanzati (SISSA-ISAS)
and INFN, Sezione di Trieste, I-34014 Trieste, Italy*



We propose and analyse a GUT-inspired version of the MSSM featuring non universal sfermion masses boundary conditions and asymptotic b - τ Yukawa coupling unification. The particle spectrum of the model gives rise to extended coannihilation processes, involving the lightest sbottom and the tau sneutrino. We study the allowed parameter space, subject to cosmological phenomenological requirements, and the details of the new coannihilations modes. Implications for direct and indirect neutralino searches, as well as for collider physics, are also briefly summarized.

1 Introduction and Motivations

The physics at the interface of cosmology and elementary particles theory is currently undergoing an exciting era. On the one side, we entered an epoch of precision measurements in the field of astrophysics and cosmology, as demonstrated by the recent results obtained by the WMAP survey¹. On the other side, the hunt for supersymmetry, which would solve a number of open fundamental questions concerning the physics beyond the standard model, is going to be boosted both by accelerator² and by direct or indirect neutralino search experiments³.

As regards the theoretical and phenomenological exploration of supersymmetric models, a number of recent experimental and observational results increasingly constrain the candidate scenarios and their parameter space. In this paper we deal with the so-called minimal supersymmetric extension of the standard model, in a *gravity mediated* SUSY breaking context⁴. In these models, the numerous *a priori* free low energy parameters are fixed, through RG running, by the values of the soft SUSY breaking masses and trilinear terms at some high energy scale M_X . The common practice is to assume that some underlying principle forces these high energy variables to some *universal* values at M_X , thus reducing the unknowns to four numbers and one sign (CMSSM).




Figure 1: The evolution of the coupling constants and of the soft SUSY breaking masses of the $\bar{\mathbf{5}}$ and $\mathbf{10}$ $SU(5)$ multiplets for a $SU(5)$ SUSY GUT theory with M_{GUT} smaller than the scale where scalar universality is assumed (here M_{Planck}).

If one further supposes that the electroweak and strong interactions ultimately unify in a single *grand unification theory* (GUT), as suggested by the running of the coupling constants, the GUT gauge group can dictate some relations among the Yukawa couplings of the high energy theory (Yukawa Unification)⁵.

We will analyse here the phenomenology of a class of GUT-inspired models, characterized by the relaxation of the hypothesis of *universality* in the sector of the fermions scalar superpartners and by partial (b - τ) Yukawa Unification (YU)⁶. These models are motivated by a minimal $SU(5)$ SUSY GUT theory in which the scale of grand unification M_{GUT} lies below the scale M_X where universality is assumed⁷. The $SU(5)$ RG running between M_X and M_{GUT} induces a non-universality in the whole scalar sector (the gaugino sector universality is instead not spoiled by the running⁷). As regards the sfermions, the multiplet structure of the GUT theory, which collects the particle content into the $\bar{\mathbf{5}}$ (down-type quark singlet and lepton doublet) and the $\mathbf{10}$ (up-type quark and lepton singlets and quark doublet) representations of $SU(5)$, implies the following parameterization for the soft SUSY breaking masses at M_{GUT} :

$$m_Q^2 = m_U^2 = m_E^2 \equiv m_{\mathbf{10}}^2 \equiv m_0^2 \quad (1)$$

$$m_L^2 = m_D^2 \equiv m_{\bar{\mathbf{5}}}^2 = K^2 m_{\mathbf{10}}^2 \quad (2)$$

In fig.1 we sketch a typical pattern for the running of $m_{\mathbf{10}}^2$ and $m_{\bar{\mathbf{5}}}^2$, compared above with the coupling constants running. Although the full GUT theory would also imply a large non-universality in the Higgs sector and $K \gtrsim 0.7$, we will resort to a simplification of the model, and let $m_{H_{1,2}}^2 = m_0^2$ in order to concentrate only on the implications of minimal non-universal sfermion masses (mNUSM). We also vary K between 0 and 1 (the latter being the fully universal case). We therefore generically expect a lighter spectrum, with respect to the universal case, for the sparticles belonging to the $\bar{\mathbf{5}}$ representation for $K < 1$.

On top of sfermion non-universality, we inherit from the original $SU(5)$ SUSY GUT the successful prediction of b - τ YU. We impose exact YU at M_{GUT} , and fix the common Yukawa coupling $h_b = h_\tau(M_{GUT})$ in order to get the right value for $m_\tau(M_Z)$, after having included running effects and the SUSY thresholds corrections at $M_{SUSY} \equiv \sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}$. The output is $m_b(M_Z)$, which is then required to lie within the properly evolved experimental range⁸.

The current upper limit on the cosmological relic density of neutralinos as cold dark matter candidates requires, in the CMSSM, efficient suppression mechanisms^{9,6}. Among these, a




Figure 2: The sbottom and sneutrino mass spectrum at $M_{1/2} = 1.1$ TeV, $\tan\beta = 38.0$ and $A_0 = 0$, for several values of m_0 , as functions of the non-universality parameter K . For the tau sneutrino, from left to right, $m_0 = 1350, 1650, 1950$ GeV, while for the sbottom $m_0 = 2850, 3300, 3750$ GeV.

recent uprise of interest has involved the so-called *coannihilation* processes⁹, occurring when the next-to-lightest SUSY particle (NLSP) mass is close to the lightest, which, for cosmological reasons, we require to be the neutralino. The main scope of the present analysis is to shed some light on the possibility that the minimal proposed deviation from universality in the sfermion soft SUSY breaking masses could activate *extended coannihilation* channels, involving sparticles belonging to the $\bar{5}$ (stau, tau sneutrino, sbottom). Further, the requirement of $b\text{-}\tau$ YU, and the fulfillment of a set of cosmo-phenomenological constraints will strongly limit the allowed $\tan\beta$ values as well as the sparticles mass range. The implications of these bounds for accelerator searches and for direct and indirect SUSY CDM detection will also be briefly discussed.

2 The Particle Spectrum

The large values of the Yukawa couplings of the third generation fermions, owing to RG running, yield lighter superpartners than the ones of the other two generations. Besides the widely discussed case of the stau⁹, the role of NLSP, in presence of non-universal sfermion boundary conditions, can be played by both the sbottom and the tau sneutrino⁶. In order to quantify this statement, we plot in fig.2 the masses of the two sparticles at various values of m_0 as functions of the parameter K . The features shown in fig.2 are traced back to the generic form of the approximate solutions to the RG equations at the electroweak scale, which can be parameterized as

$$m_{\tilde{\nu},\tilde{b}} \simeq m_0 \sqrt{a_{\tilde{\nu},\tilde{b}} + b_{\tilde{\nu},\tilde{b}} K^2}, \quad (3)$$




Figure 3: Mass isolevel curves for the tau sneutrino and for the sbottom in the $(K, m_0/M_{1/2})$ plane. The solid red and black lines indicate respectively $m_{\tilde{\nu}\tau} = 0$ and $m_{\tilde{b}} = 0$. The splitting between the isolevel curves is 10% of the neutralino mass $m_{\tilde{\chi}} \simeq 500$ GeV; $\tan\beta = 38.0$, $A_0 = 0$, $\mu < 0$.




Figure 4: (a) the neutralino relic density $\Omega_{\tilde{\chi}} h^2$ and (b) the inclusive branching ratio $BR(b \rightarrow s\gamma)$ for three benchmark K cases, corresponding respectively to $NLSP = \tilde{\tau}, \tilde{b}, \tilde{\nu}_\tau$, at $\tan\beta = 36.0$ and fixed $\Delta_{NLSP} = 10\%$.

where a and b are, to a good approximation, functions of the ratio $(m_0/M_{1/2})$ and of $\tan\beta$, and weakly depend on the trilinear coupling A_0 ⁶. In all cases we notice that for a given value of m_0 there exists a corresponding range for K where the sbottom or the tau sneutrino are quasi-degenerate with the neutralino^a (whose mass roughly depends only on $M_{1/2}$ and is therefore insensitive to K). The mass isolevel curves plotted in fig.3 highlight the plausible coannihilation regions produced by the particle spectrum of NUSM models (see sec.4). The shape of fig.3 is qualitatively unchanged by the variation of any SUSY parameter, slowly moving to the right when increasing $\tan\beta$ ⁶. We then conclude that we expect extended sfermion coannihilations for $K < 0.5$ and for relatively large values of $m_0 \gtrsim 1.5 M_{1/2}$.

3 Cosmo-Phenomenological Constraints

Supersymmetric models with conserved R parity are commonly required to fulfill two categories of phenomenological constraints. On the one side, since these models produce natural candidates for the CDM content of the Universe, in our case the neutralino, the corresponding relic abundance as well as the direct or indirect cosmic detection rates must be compatible with the current observational and experimental data. On the other side, many precision accelerator measurements can be used to test the viability of a given supersymmetric scenario. Finally, the extra theoretical constraint of b - τ YU forces the mass of the bottom quark to values to be compared with the current experimental and lattice data. Details on the numerical calculations and on the statistical analysis we adopted in this work can be found in ref.⁶.

Neutralino relic density and detection. We impose to the model the current determination of the upper limit on the CDM content of the Universe, coming from the analysis of the WMAP results¹, namely $\Omega_{CDM} h^2 = 0.1126^{+0.0081}_{-0.0091}$. The lower limit can be evaded under the hypothesis that other components contribute to the CDM besides the neutralino. The numerical computation of the neutralino relic density is performed with the updated `micrOMEGAs` code¹⁰, which includes thermally averaged exact tree-level cross-sections of all possible (co-)annihilation processes, an appropriate treatment of poles and the one-loop QCD corrections to the Higgs coupling with the fermions. Fig.4 (a) shows our results for three *benchmark* cases, respectively

^aAs regards the question of the fine-tuning of the parameter K , see the discussion in sec.6.4 of ref.⁶.




Figure 5: The cosmologically and phenomenologically allowed regions in the $(m_{\tilde{\chi}}, \Delta_{NLSP})$ plane at $\tan \beta = 38.0$, at $K = 0.1$ (a) and $K = 0.35$ (b).

at $K = 0.2, 0.4$ and 0.8 . In the latter case the coannihilation region overlaps the A -pole rapid direct annihilation funnel, extending the allowed $m_{\tilde{\chi}}$ range up to ≈ 700 GeV. The other two cases, instead, show the typical behavior $\Omega_{\tilde{\chi}} \propto m_{\tilde{\chi}}^2$, though suppressed by coannihilation phenomena.

As for the direct and indirect neutralino searches, we verified that in any case the detection rate estimates for mNUSM models lie below the *current* experimental sensitivity⁶. We find nonetheless that t -channel exchanges of lighter (for $K \ll 1$) sfermions generate higher detection rates for small values of K , falling within reach of future WIMP search experiments³.

Accelerator constraints. The most stringent bounds coming from accelerator data are found to be, in the present work, the inclusive $BR(b \rightarrow s\gamma)$ ¹¹ (see fig.4, frame (b)) and the lower limit on the lightest CP-even Higgs mass¹². They both give *lower* bounds on the mass of the neutralino $m_{\tilde{\chi}}$. As regards the SUSY contributions to the muon anomalous magnetic moment δa_μ , we chose to resort to the so-called *super-conservative* approach¹⁴, and to take as a constraint the extended 5- σ bound derived from the latest experimental and theoretical results¹³ on a_μ .

The b quark mass. In the context of b - τ YU, the SUSY corrections to the mass of the b quark constrain the allowed $\tan \beta$ range as well as the sign of μ . In fact, the sign of these corrections is given by the sign of μ , and is proportional to $\tan \beta$; since the tree level value of m_b , fixed by the common b - τ Yukawa coupling determined by m_τ , is typically higher than the upper experimental bound on m_b , the corrections are required to be *negative* (hence $\mu < 0$) and *large* (hence large $\tan \beta$). In mNUSM models⁶, the adopted top-down YU excludes $\mu > 0$ and forces, in the $\mu < 0$ case, $31 < \tan \beta < 45$.

4 Extended Sfermion Coannihilations

Minimal NUSM produce a new coannihilation branch at $K < 0.5$ (see fig.3), where the coannihilating NLSP is the tau sneutrino at low $1.5 \lesssim (m_0/M_{1/2}) \lesssim 3$, and the lightest sbottom, for $(m_0/M_{1/2}) \gtrsim 3$. We study in fig.5 the new extended sfermion coannihilations at $A_0 = 0$ and at an intermediate value of $\tan \beta = 38$. Frame (a) refers to the tau sneutrino NLSP, at $K = 0.1$, while frame (b) to the sbottom, at $K = 0.35$. We parameterize the remaining variables of the model, m_0 and $M_{1/2}$, through $m_{\tilde{\chi}}$ and $\Delta_{NLSP} \equiv \frac{m_{NLSP} - m_{\tilde{\chi}}}{m_{\tilde{\chi}}}$: via RG running, in fact, $(m_0, M_{1/2})$ uniquely determines $(m_{\tilde{\chi}}, \Delta_{NLSP})$, and vice-versa. The NLSP mass range where $\Omega_{\tilde{\chi}} \lesssim 0.13$, at a fixed given $m_{\tilde{\chi}}$, is as large as allowed by the efficiency of coannihilation

processes. We clearly notice that sbottom-neutralino coannihilations (frame (b)), which involve strong interaction processes, suppress the relic density much more effectively than in the $\tilde{\chi}-\tilde{\nu}_\tau-\tilde{\tau}$ case (frame (a)). We also find that $\tilde{\chi}-\tilde{b}$ and $\tilde{b}-\tilde{b}$ (co-)annihilations evolve, through strong interactions, into few final SM states (respectively $g-b$ and $g-g$ or $b-b^*$) and largely dominate over $\tilde{\chi}-\tilde{\chi}$ annihilations, while triple $\tilde{\chi}-\tilde{\nu}_\tau-\tilde{\tau}$ coannihilations present a wider and more complex pattern of final states (see ref. ⁶). In the left part of the figures we show the bounds dictated by the 95% C.L. exclusion regions of various accelerator constraints. Scanning the values of $\tan\beta$ we find that, for $\tan\beta \gtrsim 45$ and $\tan\beta \lesssim 31$, the cosmologically allowed region squeezes until no points simultaneously fulfill $b-\tau$ YU and the accelerator constraints.

5 Summary and Conclusions

We showed that in a $SU(5)$ SUSY GUT with $m_5 = Km_{10}$ scalar soft SUSY masses new extended sfermion coannihilation modes take place. These involve unusual NLSP coannihilating particles, such as the tau sneutrino and the sbottom. The models, to which we apply the constraint of $b-\tau$ YU, as inherited from the inspiring $SU(5)$ theory, fulfill the present phenomenological accelerator bounds as well as the recent cosmological relic density required for the neutralino as cold dark matter constituent.

As regards the detectability of the models under scrutiny, we find ⁶ that future direct WIMP search experiments ³ will be sensitive to part of the highlighted coannihilation regions, namely at low $m_{\tilde{\chi}}$. Moreover, as far as the CERN LHC pp collider reach is concerned ², we find that most of the tau-sneutrino coannihilation region will be within reach, while the sbottom coannihilation region (where $m_0 \gtrsim 3M_{1/2}$ and a heavier SUSY spectrum is cosmologically allowed), could escape detectability if $m_{\tilde{\chi}} \gtrsim 500$ GeV.

Acknowledgments

I acknowledge SISSA for financial support and the organizers of the Conference for the very pleasant and stimulating environment.

References

1. D.N. Spergel et al., [astro-ph/0302209](#); M. Limon, *Talk given at the XXXVIIIth Rencontres de Moriond, 2003*.
2. H. Baer, C. Balazs, A. Belyaev, T. Krupovnickas and X. Tata, [hep-ph/0304303](#).
3. V. Sanglard and R. Luscher, *Talks given at the XXXVIIIth Rencontres de Moriond, 2003*.
4. see e.g. A. Chamseddine, R. Arnowitt and P. Nath, *Phys. Rev. Lett.* **49** (1982) 970.
5. M. Carena, S. Pokorski and C. E. Wagner, *Nucl. Phys.* **B406** (1993) 59; D. Auto, H. Baer, C. Balazs, A. Belyaev, J. Ferrandis and X. Tata, [hep-ph/0302155](#) and references therein.
6. S. Profumo, [hep-ph/0304071](#).
7. H. Baer, M. Diaz, P. Quintana and X. Tata, *JHEP* **0004** (2000) 016.
8. H. Baer, J. Ferrandis, K. Melnikov, X. Tata, *Phys. Rev.* **D 66** (2002) 074007.
9. A. Djouadi, M. Drees and J.L. Kneur, *JHEP* **0108** (2001) 055.
10. G. Belanger, F. Boudjema, A. Pukhov and A. Semenov, *Comput. Phys. Commun.* **149** (2002) 103; [hep-ph/0210327](#).
11. R. Barate et al. (**ALEPH** Collaboration), *Phys. Lett.* **B 429** (1998) 169.
12. **ALEPH**, **DELPHI**, **L3** and **OPAL** Collaborations, [hep-ex/0107029](#).
13. A. Nyffeler and F. Gray, *Talks given at the XXXVIIIth Rencontres de Moriond, 2003*; S. Narison, [hep-ph/0303004](#).
14. S.P. Martin and J.D. Wells, *Phys. Rev.* **D 67** (2003) 015002.